



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Association of Occupational and Leisure-Time Physical Activity with Aerobic Capacity in a Working Population

Mundwiler, Jonas ; Schüpbach, Ulla ; Dieterle, Thomas ; Leuppi, Jörg Daniel ; Schmidt-Trucksäss, Arno ; Wolfer, David Paul ; Miedinger, David ; Brighenti-Zogg, Stefanie

Abstract: Objective data on the association of maximal aerobic capacity (VO₂max) with work related physical activity are sparse. Thus, it is not clear whether occupational physical activity (OPA) contributes to an increase of VO₂max. This study examined the association of VO₂max with work and non-work related physical activity in a Swiss working population.

DOI: <https://doi.org/10.1371/journal.pone.0168683>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-137481>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Mundwiler, Jonas; Schüpbach, Ulla; Dieterle, Thomas; Leuppi, Jörg Daniel; Schmidt-Trucksäss, Arno; Wolfer, David Paul; Miedinger, David; Brighenti-Zogg, Stefanie (2017). Association of Occupational and Leisure-Time Physical Activity with Aerobic Capacity in a Working Population. PLoS ONE, 12(1):1-15.
DOI: <https://doi.org/10.1371/journal.pone.0168683>


RESEARCH ARTICLE

Association of Occupational and Leisure-Time Physical Activity with Aerobic Capacity in a Working Population

Jonas Mundwiler^{1,2*}, Ulla Schüpbach¹, Thomas Dieterle^{1,3}, Jörg Daniel Leuppi^{1,3}, Arno Schmidt-Trucksäss^{2,3}, David Paul Wolfer^{4,5}, David Miedinger^{3,6}, Stefanie Brighenti-Zogg^{1,4}

1 University Clinic of Internal Medicine, Cantonal Hospital Baselland, Liestal, Switzerland, **2** Department of Sport Exercise and Health, University of Basel, Basel, Switzerland, **3** Faculty of Medicine, University of Basel, Basel, Switzerland, **4** Department of Health Sciences and Technology, ETH Zurich, Zurich, Switzerland, **5** Institute of Anatomy, University of Zurich, Zurich, Switzerland, **6** Department of Occupational Medicine, Swiss National Accident Insurance Fund (Suva), Lucerne, Switzerland

* Jonas.Mundwiler@ksbl.ch



OPEN ACCESS

Citation: Mundwiler J, Schüpbach U, Dieterle T, Leuppi JD, Schmidt-Trucksäss A, Wolfer DP, et al. (2017) Association of Occupational and Leisure-Time Physical Activity with Aerobic Capacity in a Working Population. PLoS ONE 12(1): e0168683. doi:10.1371/journal.pone.0168683

Editor: Laurent Mourot, University of France Comté, FRANCE

Received: April 11, 2016

Accepted: December 5, 2016

Published: January 3, 2017

Copyright: © 2017 Mundwiler et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This study was financially supported by an unrestricted grant of the Swiss National Accident Insurance Fund (Suva). Grant number: not applicable. URL: <http://www.suva.ch>. Recipient: JDL. The funder (Suva) provided support in the form of a project grant that covered salaries for the involved personnel at the Cantonal Hospital Baselland Liestal as well as expenses for

Abstract

Introduction

Objective data on the association of maximal aerobic capacity (VO_{2max}) with work related physical activity are sparse. Thus, it is not clear whether occupational physical activity (OPA) contributes to an increase of VO_{2max} . This study examined the association of VO_{2max} with work and non-work related physical activity in a Swiss working population.

Methods

In this cross-sectional study, a total of 337 healthy and full-time employed adults were recruited. Demographic data, height, weight and BMI were recorded in all subjects. Participants were classified into nine occupational categories (ISCO-88) and merged into three groups with low, moderate, and high OPA. Physical activity was objectively measured by the SenseWear Mini Armband on seven consecutive days (23 hours per day). Participants were regarded as sufficiently active when accumulating ≥ 30 min of moderate-to-vigorous physical activity per day. VO_{2max} was evaluated using the multistage 20-meter shuttle run test.

Results

Data of 303 participants were considered for analysis (63% male, age 33 yrs, SD 12). Multiple linear regression analysis (adjusted $R^2 = 0.69$) revealed significant positive associations of VO_{2max} with leisure-time physical activity (LTPA) at vigorous intensity ($\beta = 0.212$) and sufficient moderate-to-vigorous physical activity ($\beta = 0.100$) on workdays. Female gender ($\beta = -0.622$), age ($\beta = -0.264$), BMI ($\beta = -0.220$), the ratio of maximum to resting heart rate ($\beta = 0.192$), occupational group (low vs. high OPA, $\beta = -0.141$), and smoking ($\beta = -0.133$) were also identified as independent predictors of VO_{2max} .

infrastructure and conducting the study, but did not have any role in the study design, data collection and analysis, decision to publish, or preparation of this manuscript. The participation of DM in the study was independent from the project grant and is covered by the contractual agreement between Suva and DM that permits active participation in independently conducted research projects.

Competing Interests: DM is an employee of the funding organization (Suva). Beside his work as a consultant at Suva, he is involved in independent research projects at the Cantonal Hospital Baselland Liestal and the University of Basel. The funder (Suva) did not have any role in the study design, data collection and analysis, decision to publish, or preparation of this manuscript. Furthermore, this does not alter our adherence to PLOS ONE policies on sharing data and materials. The other authors have declared that no competing interests exist.

Conclusions

The present results suggest that VO_{2max} is positively associated with LTPA, but not with OPA on workdays. This finding emphasizes the need for employees to engage in sufficient high-intensity physical activity in recreation for maintaining or improving VO_{2max} with regard to health benefits.

Introduction

VO_{2max} is defined as the maximum rate of oxygen consumption. It is generally accepted as an appropriate measure of the functional capacity of the cardiorespiratory system and is commonly interpreted as an index of cardiorespiratory fitness [1]. Previous studies found that poor cardiorespiratory fitness was a risk factor for various diseases such as hypertension, stroke, type 2 diabetes, and metabolic syndrome [2, 3]. Other studies reported that a low VO_{2max} was associated with all-cause mortality and mortality from cardiovascular disease [4].

VO_{2max} is determined by genetic factors, age, gender, as well as physical activity, body fat, smoking, and medical conditions, i.e. metabolic syndrome and diabetes [5]. VO_{2max} decreases with age with an average rate of decline of about 1% per year or 10% per decade after the age of 25 [6, 7]. Recent data suggest that the ability of an individual to increase VO_{2max} is genetically determined. Each individual disposes of a predetermined genetic window, and can vary the amount of VO_{2max} with exercise training or detraining within that window [8]. VO_{2max} values range from about 10 ml/kg/min in severely ill cardiac patients to 80–90 ml/kg/min in world-class runners and cross-country skiers [9]. VO_{2max} may be substantially increased in response to endurance training [8].

For decades, governmental and non-governmental agencies have promoted physical activity for individuals' health benefits. The World Health Organization recommends that adults between 18 and 64 years should engage in ≥ 30 min of at least moderate-intensity physical activity on most days of the week. Aerobic activity should be performed in bouts of ≥ 10 min duration across different domains such as work, leisure-time, transport, domestic and garden [10]. Since most people in full-time employment spend one third or more of the day at work, it is conceivable that occupational physical activity (OPA) may contribute to a large extent to total daily activity. While leisure-time physical activity (LTPA) is well known to be positively associated with VO_{2max} [8], only few data are available on the relationship between OPA and VO_{2max} in employees. With regards to LTPA, Ong & Sothy [11] found that regularly exercising men in sedentary occupations had a significantly higher mean VO_{2max} than non-regularly exercising counterparts. In contrast, the potential positive effects of OPA on VO_{2max} are less well investigated [12]. However, available data suggest that OPA, independent of formal exercise programs, may positively affect VO_{2max} [13].

Previous studies measuring physical activity in employees used pedometers and accelerometers in combination with self-reported questionnaires. Self-reported questionnaires are the most used method for physical activity assessment [14]. However, validation studies comparing self-reported estimates to the gold standard method Doubly-Labeled-Water (DLW) are inconsistent [15, 16]. Due to accuracy, ability to capture large amounts of data and ease of administration, accelerometers are widely used today [17]. However, they are not accurate in estimating physical activity during activities involving uphill and downhill walking or when carrying heavy load while walking [18]. Pedometers directly observe duration of activities, but do not record intensity and frequency of physical activity or purely upper body movements

[19]. Armband devices, such as the SenseWear Mini Armband, integrate motion and heat-related sensors. This dual measurement strategy is more sensitive for assessing energy expenditure associated with complex and non-ambulatory activities, such as carrying heavy load while walking [19]. Furthermore, this method ensures a sensitive determination of acceleration provoked by muscle power or externally by a vehicle or gravitation [20].

Due to methodological limitations and lack of studies, potential associations of objective physical activity and VO_{2max} in employees still need to be clarified by further investigations. Furthermore, with regards to the development of evidence-based activity recommendations it is important to know, whether OPA contributes to an improvement of VO_{2max} . Therefore, the objective of this study was to analyse the relationship between VO_{2max} and work and non-work related physical activity as measured by the SenseWear Mini Armband in a Swiss working population. The secondary objective was to evaluate the effect of demographic factors on VO_{2max} including gender, age, body mass index (BMI), and smoking.

Materials and Methods

Study Participants

From May 17, 2013 (first participant in) to February 11, 2015 (last participant in), a total of 337 healthy and at minimum 80% employed adult workers from various companies of the Basel region, Switzerland were recruited. Exclusion criteria were insufficient knowledge of the German language, movement restrictions as well as diseases and accidents within the last three months that affected productivity at the workplace. Furthermore, individuals, who had to comply with specific security regulations, and night shift workers could not take part in this study because of their altered sleep, eating and physical activity behaviour. This investigation has been conducted according to the Declaration of Helsinki and was approved by the local ethics committee “Ethikkommission Nordwest- und Zentralschweiz” (EKNZ, 260/12) on December 21, 2012. Written informed consent was obtained from all study participants prior to study entry.

Study Design and Procedures

In this cross-sectional study, the aim was to recruit an equal distribution of subjects across different occupational groups. A permit from leading persons of miscellaneous companies was requested to receive contact details for potentially recruitable employees, who were then informed and asked for study participation by phone or by e-mail. The selected companies included medium sized corporations from the public sector (e. g. hospitals) as well as small sized private firms (e. g. construction companies). At the study visit, height and weight were reliably measured. Height was assessed without shoes by a medical measuring stick to the nearest mm (model Seca 217, measurement range: 20 to 205 cm, Seca AG, Reinach, Switzerland). The measurement of weight was performed on subjects in light clothing without shoes by a medical scale with an accuracy of 0.1 kg (model Seca 877, load capacity: 200 kg, Seca AG, Reinach, Switzerland). BMI was calculated from measured height and weight ($BMI = \text{weight} / \text{height}^2$ [kg/m^2]). Subjects with a BMI of $25 \text{ kg}/\text{m}^2$ or more were classified as overweight, and those with a BMI of $30 \text{ kg}/\text{m}^2$ or more as obese [21]. In addition, a variety of personal and job-related factors were recorded by a generic questionnaire, such as age, gender, nationality, marital status, smoking status, alcohol consumption, highest education, current profession, daily working hours, working time model, medication, psychotherapy, illnesses and accidents within the last three months. The reported professions were classified into nine categories based on the International Standard Classification of Occupations 1988 (ISCO-88) [22]. Participants were then merged into three groups with low (managers, scientists, office workers),

moderate (technicians, service workers, machine operators), and high (agricultural workers, craftsmen, labourers) OPA [23]. Prior to the observation period, subjects performed a 20-meter shuttle run test in order to determine VO_{2max} . During the subsequent week, participants were instructed to wear the SenseWear Mini Armband on seven consecutive days in order to objectively measure daily physical activity.

Measurements

20-meter shuttle run test. The multistage 20-meter shuttle run test is a common endurance fitness test to evaluate the maximal aerobic capacity of healthy adults. It is simple in use, economical and large groups can be tested simultaneously. Validity of the one-minute stage version of the 20-meter shuttle run to predict VO_{2max} in adults was established by Léger & Gadoury [24], who compared the maximal shuttle run speed to VO_{2max} attained during a multistage treadmill test ($r = 0.90$). Test-retest reliability was found to be very high ($r = 0.95$) in healthy adults [23].

This test was conducted on a flat, non-slip surface. Participants were instructed to run back and forth between two lines, which were 20-meters apart, with a running velocity determined by audio signals [23]. Starting speed was 8.5 km/h and every minute (stage), speed was increased by 0.5 km/h until the subject could no longer keep the pace and did not reach the lines in time twice in a row [23]. The test result corresponded to the number of reached stages. According to a validated table [25], this score was used to predict VO_{2max} , which could be compared to age-dependent normative data for males and females.

Resting heart rate (HR_{rest}), systolic and diastolic blood pressure were assessed prior to testing after 10 minutes at rest in a sitting position in a quiet environment. Heart rate was continuously recorded during the test up to maximum frequency (HR_{max}) and recovery pulse ($HR_{recovery}$), systolic and diastolic blood pressure were measured two minutes after the end of the test (in a sitting position in a quiet environment). The ratios of HR_{max} -to- HR_{rest} and HR_{max} -to- $HR_{recovery}$ were calculated as (HR_{max}/HR_{rest}) and ($HR_{max}/HR_{recovery}$) [26], respectively. Hypertension was defined as a blood pressure of $\geq 140/90$ mmHg. Four participants did not perform the 20-meter shuttle run due to a resting systolic blood pressure > 180 mmHg. They were pairwise excluded from the corresponding analyses.

SenseWear mini armband. The SenseWear Mini Armband (Model MF-SW) is a small, lightweight and wireless multisensory activity monitor developed by BodyMedia Inc., Pittsburgh, Pennsylvania, USA (now Jawbone Inc., San Francisco, California, USA), which integrates motion data from a three-axis accelerometer along with other sensors such as heat flux, skin temperature and galvanic skin response. Validity of the SenseWear Mini Armband was established by Johannsen *et al.* [20] comparing energy expenditure estimates of the SenseWear Mini Armband against the criterion method DLW in healthy adults. The Armband showed a high intraclass correlation with DLW ($r = 0.85$) and a low absolute error rate (8%, SD 7%) [20].

Subjects were instructed to wear the SenseWear Mini Armband on the upper left arm (triceps area) for seven consecutive days, including while sleeping, with the exception of the time spent on personal hygiene. The first and the last incomplete measurement day, including the study visits, were not taken into account. Therefore, the investigated measurement period was five days, which had to consist of at least three workdays to be included in the analysis [27]. A day was considered as a whole workday, if participants worked cumulatively ≥ 6 h, and as a half workday in case of ≥ 3 to < 6 h. Days with < 3 working h were regarded as non-working days. Measurement periods of < 22 h per day or < 12 h during wake time were excluded from analysis [28]. Information about workdays and non-working days as well as work-time and

Table 1. List of SenseWear physical activity variables and their definition.

Variable	Definition
MPA	Sum of moderate (one-minute intervals) physical activity (≥ 3 to < 6 METs) during work-time and during leisure-time on workdays / on non-working days.
HPA	Sum of high (one-minute intervals) physical activity (≥ 6 to < 9 METs) during work-time and during leisure-time on workdays / on non-working days.
VHPA	Sum of very high (one-minute intervals) physical activity (≥ 9 METs) during work-time and during leisure-time on workdays / on non-working days.
VPA	Sum of vigorous (one-minute intervals) physical activity (≥ 6 METs) (combination of HPA and VHPA, produced for the regression analysis).
MVPA	Sum of sufficient (≥ 30 min) / insufficient (< 30 min) moderate-to-vigorous (≥ 3 METs) physical activity (bouts of ≥ 10 min) on workdays / non-working days.

doi:10.1371/journal.pone.0168683.t001

leisure-time on workdays was obtained from diaries participants filled in during the measurement period.

Calculation of physical activity scores. The physiological data, collected by the arm-band's sensors, were processed by specific algorithms available in the SenseWear software (BodyMedia, professional software V.7.0, algorithm V.2.2.4). Patients' average daily number of steps, active energy expenditure (AEE), physical activity level in metabolic equivalents of task (METs) and physical activity duration at different intensities were examined. One MET corresponds to 3.5 ml/min/kg VO_2 [29]. For all variables, average values were calculated separately for workdays and non-working days as well as for work-time and leisure-time on workdays. The amount of physical activity (min/day) at a certain intensity level was calculated in two ways (Table 1). First, one-minute intervals in which the intensity reached the following MET thresholds were summed up: moderate physical activity (MPA) ≥ 3 to < 6 METs, high physical activity (HPA) ≥ 6 to < 9 METs, and very high physical activity (VHPA) ≥ 9 METs. As participants performed only little physical activity in the VHPA range, this variable has been combined with HPA for a more representative measure with respect to the regression analysis (combined variable: vigorous physical activity (VPA)). Second, because current guidelines suggest accumulating physical activity bouts of ≥ 10 min [29], this was considered in the calculation of moderate-to-vigorous physical activity (MVPA) ≥ 3 METs. Thus, we were able to investigate whether participants fulfilled the recommendation of MVPA of ≥ 30 min per day calculated from bouts ≥ 10 min on workdays and non-working days [30].

Statistical Analysis

Data were analyzed using IBM SPSS Statistics (version 22.0). Significance was set at the 5% level. The Shapiro-Wilk test was used to test whether data were normally distributed. Data are presented as mean and standard deviation (SD) or number and percentage. To analyse differences across gender and physical activity categories, mean comparisons were performed using Student's T-test or Mann-Whitney test, if appropriate. Categorical data were analyzed with Chi-Square test. To identify potential predictors of VO_{2max} , a multiple linear regression analysis was performed using the backward stepwise method. VO_{2max} was considered as dependent variable. Independent variables were age, gender, BMI, AEE / MPA / VPA / Steps during work-time, leisure-time and on non-working days, MVPA on workdays and non-working days (insufficient vs. sufficient), HR_{max} -to- HR_{rest} , smoking (never smokers vs. current smokers / never smokers vs. ex-smokers), and occupational group (group 1 vs. group 2 / group 1 vs. group 3). Validity of the regression model was established by checking essential assumptions. Checks for collinearity were made using tolerance and variance inflation factor (VIF) for each

of the independent variables calculated by means of the collinearity diagnostics in IBM SPSS Statistics. AEE measured by the SenseWear Mini Armband was subject to power calculation. Assuming a sample size of 100 subjects in each occupational group, there is a power of >90% to detect a mean difference of 500 kcal between any of these groups. This calculation was based on the assumption of a within group SD of 730 kcal and on a two-sided significance level of 5% [31].

Results

Subjects' Characteristics

Of the 337 recruited subjects 303 were considered for analysis. Descriptive data for total ($n = 303$), male ($n = 190$, 63%), and female ($n = 113$, 37%) subjects are presented in Table 2. 31% of subjects ($n = 95$) were found to be overweight and 7% ($n = 21$) were obese. Age ranged from 18 to 61 years and did not differ significantly between sexes. A higher percentage of males compared to females were current smokers, while more women than men were ex-smokers. BMI, HR_{max} , $HR_{recovery}$, and systolic blood pressure were significantly increased in male subjects. In contrast, only HR_{rest} was higher in females. Diastolic blood pressure was about the same in men and woman. More than half of the men suffered from hypertension, whereas in woman less than one-third was affected. Gender distribution was more or less balanced in group 1 (low OPA) and group 2 (moderate OPA), while only 4% of the investigated females were represented in group 3 (high OPA).

Thirty-four subjects (10%) have worn the SWMA on less than three workdays and were therefore excluded from the entire analysis. Reasons for non-wearing or non-evaluation were:

Table 2. Characteristics of study subjects.

Variables	Total ($n = 303$)	Male ($n = 190$)	Female ($n = 113$)
	Mean (SD) or N (%)	Mean (SD) or N (%)	Mean (SD) or N (%)
Age [yrs]	33 (12)	33 (13)	35 (12)
BMI [kg/m^2]	24 (3)	25 (3) ***	23 (4)
Current smokers	64 (21%)	46 (24%)	18 (16%)
Ex-smokers	61 (20%)	32 (17%)	28 (25%)
HR_{rest}	72 (13)	70 (13)*	74 (13)
HR_{max}	183 (15)	187 (14) ***	178 (15)
$HR_{recovery}$	106 (15)	108 (14) **	102 (16)
Sys BP _{rest} [mmHg]	135 (15)	140 (12)**	128 (18)
Dia BP _{rest} [mmHg]	82 (10)	82 (10)	80 (11)
Hypertension ($\geq 140/90$ mmHg)	130 (43%)	104 (55%)	26 (23%)
History of Diabetes	0 (0%)	0 (0%)	0 (0%)
History of CAD	1 (0%)	1 (0%)	0 (0%)
Group 1 (low OPA)	101 (33%)	55 (29%)	46 (41%)
Group 2 (moderate OPA)	102 (34%)	40 (21%)	62 (55%)
Group 3 (high OPA)	100 (33%)	95 (50%) **	5 (4%)

BMI, body mass index; CAD, coronary artery disease, Dia BP_{rest}, diastolic blood pressure at rest; HR_{max} , maximum heart rate; $HR_{recovery}$, recovery pulse (measured two minutes after the end of the 20-meter shuttle run test); HR_{rest} , heart rate at rest; OPA, occupational physical activity; SD, standard deviation; Sys BP_{rest}, systolic blood pressure at rest; VO_{2max} , maximal oxygen consumption during multistage 20-meter shuttle run test.

* $p < 0.05$,

** $p < 0.01$,

*** $p < 0.001$ (two-tailed) vs. females.

doi:10.1371/journal.pone.0168683.t002

technical problems ($n = 4$), illness during observation period ($n = 2$), no paid occupation ($n = 5$), no interest ($n = 15$), loss of the armband ($n = 4$), skin irritations ($n = 2$) or sleep problems ($n = 2$). Another 24 individuals had missing SWMA data on non-working days due to more workdays during observation period and were pairwise excluded from the corresponding analyses.

Physical Activity Data

Table 3 illustrates VO_{2max} and objective physical activity measured by the SenseWear Mini Armband according to gender. Males had a significantly higher VO_{2max} than females. Overall, most physical activity was performed within the moderate-intensity range. Furthermore, activity levels at all intensities, AEE, and the number of daily steps were higher in males compared to females. Activity parameters differed significantly between sexes on workdays during work- and leisure-time (with the exception of VHPA during leisure-time), while no gender-dependent differences in physical activity were found on non-working days (with the exception of AEE). Moreover, men showed higher physical activity levels on workdays (work-time and leisure-time added) than on non-working days. In women, this was the case for MPA, AEE, and steps, whereas it was the contrary for HPA and VHPA.

VO_{2max} across Categories of Physical Activity

On workdays, VO_{2max} was significantly higher in participants, who were sufficiently active (≥ 30 min MVPA/day, 42 ml/min/kg, SD 9, $n = 251$, 84%) compared to those, who were insufficiently active (< 30 min MVPA/day, 31 ml/min/kg, SD 7, $n = 48$, 16%) ($p < 0.01$). However,

Table 3. Physical activity data, measured with objective methodology, SenseWear Mini Armband.

Variables		Total ($n = 303$)	Male ($n = 190$)	Female ($n = 113$)
		Mean (SD)	Mean (SD)	Mean (SD)
VO_{2max}	[ml/kg/min]	40 (10)	45 (8) ***	33 (7)
MPA	Work-time [min/day]	151 (131)	190 (143) ***	85 (70)
	Leisure-time [min/day]	107 (50)	113 (52) *	98 (44)
	Non-working day [min/day]	178 (105)	187 (110)	164 (94)
HPA	Work-time [min/day]	8 (14)	11 (16) ***	2 (6)
	Leisure-time [min/day]	9 (11)	10 (11) **	7 (11)
	Non-working day [min/day]	12 (18)	13 (21)	9 (13)
VHPA	Work-time [min/day]	0 (1)	0 (2) **	0 (0)
	Leisure-time [min/day]	2 (5)	2 (5)	1 (4)
	Non-working day [min/day]	2 (7)	2 (8)	2 (6)
AEE	Work-time [kcal/day]	795 (816)	1068 (905) ***	337 (273)
	Leisure-time [kcal/day]	609 (321)	700 (341) ***	456 (211)
	Non-working day [kcal/day]	950 (632)	1089 (706) ***	724 (398)
Daily steps	Work-time	6521 (3888)	7511 (4292) ***	4855 (2285)
	Leisure-time	5638 (2592)	5470 (2727) *	5919 (2332)
	Non-working day	8997 (6190)	9187 (7321)	8685 (3673)

AEE, active energy expenditure; MPA / HPA / VHPA, physical activity duration at moderate (3–6 METs) / high (6–9 METs) / very high (≥ 9 METs) intensity; SD, standard deviation; VO_{2max} , maximal oxygen consumption during multistage 20-meter shuttle run test.

* $p < 0.05$,

** $p < 0.01$,

*** $p < 0.001$ (two-tailed) vs. females.

doi:10.1371/journal.pone.0168683.t003

Table 4. VO_{2max} reference values according to age and gender.

		VO_{2max} [ml/kg/min]	Decrease over decade	VO_{2max} [ml/kg/min] ranges for non-athletes according to Kenney <i>et al.</i> [8]
Age categories	N (%)	Mean (SD)	%	Mean (reference ranges)
Total:	303			
18–29 yrs	131 (43)	43 (9)		
30–39 yrs	77 (26)	41 (9)	- 5%	
40–49 yrs	52 (17)	39 (8)	- 5%	
50–61 yrs	43 (14)	33 (8)	-15%	
Male:	190			
18–29 yrs	91 (48)	47 (7)		48 (43–52)
30–39 yrs	40 (21)	45 (9)	- 4%	44 (39–48)
40–49 yrs	36 (19)	42 (7)	- 7%	40 (36–44)
50–61 yrs	23 (12)	38 (6)	- 10%	38 (34–41)
Female:	113			
18–29 yrs	40 (35)	34 (7)		38 (33–42)
30–39 yrs	37 (33)	36 (6)	+ 6%	34 (30–38)
40–49 yrs	16 (14)	31 (5)	- 14%	31 (26–35)
50–61 yrs	20 (18)	26 (4)	- 16%	29 (24–33)

SD, standard deviation; VO_{2max} , maximal oxygen consumption during multistage 20-meter shuttle run test.

doi:10.1371/journal.pone.0168683.t004

on non-working days, the difference between sufficiently (41 ml/min/kg, SD 9, $n = 208$, 75%) and insufficiently (37 ml/min/kg, SD 10, $n = 71$, 25%) active participants was not significant ($p = 0.129$).

Regarding the combination of physical activity categories on workdays and non-working days, VO_{2max} was found to be lowest in the category, which did not fulfil the activity recommendations on both, workdays and non-working days (30 ml/min/kg), slightly increased in the category, which fulfilled the recommendations on non-working days, but not on workdays (34 ml/min/kg), clearly higher in those, who were sufficiently active on workdays, but not on non-working days (41 ml/min/kg) and highest in the category, which fulfilled the recommendations on both, workdays and non-working days (42 ml/min/kg).

VO_{2max} Reference Values

Average VO_{2max} values stratified by age and gender are given in Table 4. They were found to be well within the representative reference ranges of VO_{2max} for non-athletes provided by Kenney *et al.* [8].

Independent Predictors of VO_{2max}

Results of the backward stepwise multiple linear regression analysis with VO_{2max} as dependent variable are presented in Table 5. Adjusted R^2 of the model was high explaining 69% of variance in VO_{2max} . In decreasing order, female gender, age, BMI, VPA in leisure-time, HR_{max} -to- HR_{rest} occupational group 3 (vs. group 1), smoking, as well as sufficient MVPA on workdays contributed significantly to the model. In contrast, AEE / MPA / VPA / Steps during work-time, MPA / AEE / Steps in leisure-time, AEE / MPA / VPA / Steps and sufficient MVPA on non-working days, ex-smoking, occupational group 2 (vs. group 1) and HR_{max} -to- $HR_{recovery}$ were not found to be significant predictors of VO_{2max} . Based on the results of the multiple

Table 5. Backward stepwise multiple linear regression analysis with VO_{2max} as dependent variable.

Model: n = 273, Adjusted R^2 = 0.69	B	SE B	β	p-value
Constant	63.006	4.882		<0.001
Gender (male vs. female)	-11.902	0.811	-0.622	<0.001
Age [yrs]	-0.197	0.029	-0.264	<0.001
BMI [kg/m ²]	-0.602	0.112	-0.220	<0.001
VPA leisure-time [min/day]	0.155	0.027	0.212	<0.001
HR _{max} -to-HR _{rest}	3.790	0.720	0.192	<0.001
Occupational group (group 1 vs. group 3)	-2.788	0.842	-0.141	<0.001
Smoking (never vs. current smokers)	-3.061	0.811	-0.133	<0.001
MVPA workday (insufficient vs. sufficient)	2.506	1.001	0.100	0.013

B, unstandardized regression coefficient; β , standardized beta coefficient; BMI, body mass index; HR_{max}-to-HR_{rest}, ratio of maximum to resting heart rate; MVPA workday (insufficient vs. sufficient), comparison of insufficient (<30 min per day) to sufficient (\geq 30 min per day) moderate-to-vigorous physical activity calculated from bouts of \geq 10 min on workdays; Occupational group (group 1 vs. group 3), comparison of group 1 with low occupational physical activity to group 3 with high occupational physical activity; SE, standard error; VO_{2max} , maximal oxygen consumption during multistage 20-meter shuttle run test; VPA, physical activity duration at vigorous (\geq 6 METs) intensity. Excluded variables were: AEE / MPA / VPA / Steps during work-time, AEE / MPA / Steps in leisure-time, AEE / MPA / VPA / Steps on non-working days, ex-smokers (vs. never smokers), occupational group 2 (vs. group 1), HR_{max}-to-HR_{recovery}, MVPA non-working days (insufficient vs. sufficient).

doi:10.1371/journal.pone.0168683.t005

linear regression analysis, this study has generated the following prediction equation:

$$\begin{aligned}
 VO_{2max}[\text{ml/kg/min}] = & 63.006 - (11.902 \times \text{gender; men} = 1, \text{ women} = 2) - (0.197 \times \text{age [yrs]}) \\
 & - (0.602 \times \text{BMI [kg/m}^2]) + (0.155 \times \text{VPA leisure} - \text{time [min/day]}) + (3.790 \times \text{HR}_{\text{max}} - \text{to} - \text{HR}_{\text{rest}}) \\
 & - (2.788 \times \text{occupational group; group 1} = 0, \text{ group 3} = 1) - (3.061 \times \text{smoking; never smokers} = 0, \\
 & \text{current smokers} = 1) + (2.506 \times \text{MVPA workday; insufficient} = 1, \text{ sufficient} = 2).
 \end{aligned}$$

Discussion

Main Findings

This cross-sectional study found that men had higher levels of both, physical activity and VO_{2max} than women. In general, physical activity was more common during workdays than during non-working days, especially among men. In addition, on workdays, mean VO_{2max} was significantly higher in participants, who fulfilled the global recommendations on physical activity compared to insufficiently active counterparts. Multiple linear regression analysis with VO_{2max} as dependent variable showed significant positive associations of VO_{2max} with LTPA at high intensities and with sufficient MVPA on workdays, but not with OPA at any intensity. Female gender, age, BMI, the ratio of maximum to resting heart rate, occupational group 3 (vs. group 1), and smoking appeared as additional independent predictors of VO_{2max} .

Predictors of VO_{2max}

The present results are in line with existing knowledge that females usually present lower levels of physical activity and VO_{2max} than males. The regression model with objective SenseWear activity data has shown that female gender is the strongest predictor of VO_{2max} . The gender difference in VO_{2max} can partly be explained by a higher percentage of body fat and a lower hemoglobin concentration in females compared to males. Furthermore, men have a

higher proportion of muscle mass and thus more mitochondria than females [8]. The decline of VO_{2max} with age in the present study is comparable to previous studies of Jackson *et al.* [6, 7]. In this study, participants younger than 50 years had a mean decline <10% per decade and those older than 50 years a decline >10% per decade. As shown by previous studies, measured VO_{2max} presented a significant negative association with BMI. The ratio of HR_{max} -to- HR_{rest} contributed positively to the model, which is in line with a study of well-trained men by Uth *et al.* [26], who observed highly significant correlations between measured VO_{2max} and the Heart Rate Ratio Method. Consistent with previous investigations, measured VO_{2max} was negatively associated with smoking [32]. In this study, only VPA in leisure-time contributed significantly to the regression model and showed a positive association with VO_{2max} . Kenney *et al.* [8] stated that the higher the initial state of fitness, the smaller is the relative improvement for the same volume of training. Subjects with a mean baseline VO_{2max} of 40–51 ml/kg/min required an intensity of at least 45% of oxygen uptake reserve (\pm 4.7–6.1 METs) to improve VO_{2max} [33]. Since VO_{2max} values of a large proportion of the present sample were within this range (mean VO_{2max} = 40 ml/kg/min, SD 10), this study could confirm the statement, even if the threshold of 4.7–6.1 METs still falls within the range of MPA, but at the upper limit. Contrary to expectation, OPA at any intensity was not positively associated with VO_{2max} . A possible reason could be that OPA despite its long duration is more intermittent and not as effective as LTPA, which in general is planned, structured, often short, of high intensity and very efficient [29]. However, the present results are contradictory to previous findings. Hammermeister *et al.* [34] and Jang *et al.* [12] suggested that OPA could have a significant effect on VO_{2max} . Hirai *et al.* [35] also found a significant relationship between work form and VO_{2max} in male workers. The reason that they found positive associations between OPA and VO_{2max} might be that they used simple self-reported questionnaires for assessing OPA. Miller & Brown [36] stated that measuring work-related activity is problematic, because the intermittent and unstructured nature of most work-related activities makes self-report difficult. The finding that subjects in occupational group 3 with high OPA had a lower VO_{2max} than those in group 1 with low OPA could be explained by the lack of motivation to engage in high-intensity sports and exercises after a tiring work-shift, since only VPA in leisure-time was found to be predictive for VO_{2max} .

Generalizability of Results

The present study included a wide range of manual and non-manual employees and represented a typical cross-section of the Swiss working population. Mean BMI, percentage of overweight, obesity, as well as gender distribution were comparable to the Swiss working population in 2014 [37]. Furthermore, VO_{2max} values in this study were in high agreement with a previous population-based study in US employees [38]. This applies to mean total VO_{2max} as well as mean VO_{2max} in men and women. Physical activity data measured by the SenseWear Mini Armband were comparable to a recent study that objectively measured physical activity in 9554 Finnish employees with the Firstbeat Bodyguard device (dedicated device for beat-by-beat 24 h heart rate and heart rate variability measurement) [39]. In both studies, most physical activity was performed within the moderate-intensity range, and activity levels at all intensities were higher in males compared to females. Contradictory to this study was that on non-working days men accumulated more physical activity at all intensities than on workdays. Concerning women, the present results were consistent with their findings. In both studies, women showed more MPA on non-working days compared to workdays, while it was the contrary for HPA and VHPA.

Findings in Relation to Physical Activity Recommendations

In this study, 84% and 75% of participants fulfilled the activity recommendations of ≥ 30 min MVPA per day on workdays and non-working days, respectively. These findings were slightly higher compared to a survey in the Swiss population from 2012, where 72% were sufficiently active in their leisure-time [40]. However, in the survey a self-reported questionnaire was used to assess physical activity and only LTPA was considered. This might explain the higher percentage of participants fulfilling the activity recommendations in the present study compared to the Swiss survey of 2012. An explanation for the findings that LTPA on workdays was predictive for VO_{2max} and participants being sufficiently active on workdays presented a higher VO_{2max} , could be that on workdays employees only have a little time window for exercising outside work, which increases the density of activities. In contrast, on non-working days physical activity may be more unstructured and less efficient because of the extended time availability (lower density of physical activity).

Clinical Implications

The potential health benefits of good cardiorespiratory fitness (high VO_{2max} value) are well known. It reduces the risk of various diseases in the general population. This investigation has shown that OPA does not contribute to an improvement of VO_{2max} . In contrast, to maintain or improve VO_{2max} , intensive physical exercise in leisure-time in the range of high-to-very high intensity is required, as for example athletic cycling, soccer, martial arts, squash, inline skating or aerobics [41]. A low VO_{2max} and insufficient physical activity have a strong impact on individuals' wellbeing and all-cause mortality [2, 3]. Based on the present findings, it may be recommended to implement an attractive and intensive sports program at the workplace, such as lunch-time or after-work exercise in order to improve the overall health in the working population.

Strengths and Limitations of the Study

The present study had several strengths. The study sample included a wide range of manual and non-manual employees and represented a typical cross-section of the Swiss working population. Furthermore, the measurement of physical activity and VO_{2max} was conducted with objective instruments. The SenseWear Mini Armband promises an accurate assessment of physical activity under non-ambulatory conditions. The inclusion of thermal- and perspiration-related sensors in addition to the three-axis accelerometer provides a way to detect subtle increases in physical activity associated with low-intensity activities [20]. Furthermore, the detection of non-wearing, resting and sleep time allow for more confidence in data consistency. However, there are also some limitations associated with the Sense Wear Mini Armband. The device has been shown to underestimate physical activity at very high intensities (>10 METs) [42]. This may explain why in the present study only few subjects had activities above this intensity threshold (VHPA). Though, the relationship between VO_{2max} and vigorous LTPA is expected to be even stronger with reliable measurements of VHPA. Moreover, the Sense Wear Mini Armband was found to underestimate activities involving purely lower extremities, such as cycling, because of its wearing position on the upper arm [43]. In addition, it is not waterproof and lacks to detect water-based activities. A strength of this study were the strict exclusion criteria for recording days (e. g. measurement periods of <22 h per day or <12 h during wake time were excluded from analysis). Thus, the recordings covered well typical workdays and non-working days. Nevertheless, the present study also had some weaknesses. Since only 4% of women participated in group 3 (high OPA), there is a need for further investigations to focus on females in this subgroup. Another limitation was that VO_{2max} values were

not directly measured, but predicted from the score, which participants reached in the 20-meter shuttle run test. For direct measurement of VO_{2max} , spiroergometry is considered to be the gold standard [44]. However, spiroergometry is labour-intensive, requires trained staff and is therefore not feasible for assessing VO_{2max} in large populations [45]. A recent study confirmed the validity of the 20-meter shuttle run test and concluded that it can accurately predict VO_{2max} in healthy adults [46]. The results revealed significant correlations between the number of shuttles in the 20-meter shuttle run test and directly measured VO_{2max} ($r = 0.87$, $p < 0.05$) as well as the velocity at which VO_{2max} occurred ($r = 0.93$, $p < 0.05$) [46]. The present study was cross-sectional in nature, which provided a snapshot of the relation between physical activity and VO_{2max} . However, cause-effect-relationships are difficult to ascertain. To overcome this limitation in future, longitudinal intervention studies are required.

Conclusions

The key finding of this cross-sectional investigation is that VO_{2max} was not positively associated with OPA in a representative cohort of healthy Swiss employees, when objectively measured with the SenseWear Mini Armband. In accordance with existing knowledge, VO_{2max} showed a positive association with LTPA at vigorous intensity and with sufficient MVPA on workdays. Furthermore, female gender, age, BMI, the ratio of maximum to resting heart rate, occupational group 3 (vs. group 1), and smoking were identified as independent predictors of VO_{2max} . These findings provide the basis and therefore will be of great value for developing evidence-based strategies to improve cardiorespiratory fitness in the working population.

Supporting Information

S1 Table. Professions of study subjects, stratified according to occupational group and category.

(DOCX)

S1 Dataset. Excel file of personal and job-related factors.

(XLSX)

S2 Dataset. Excel file of body measurements and 20-meter shuttle run.

(XLSX)

S3 Dataset. Excel file of SenseWear Mini Armband activity parameters.

(XLSX)

S1 Protocol. Study protocol approved by the local ethics committee.

(DOC)

Acknowledgments

The authors of the present study would like to thank all employees for their study participation. In addition, the authors express their appreciation to “Gewerblich-industrielle Berufsfachschule Liestal” and “Kaserne Liestal” for their technical and equipment support.

Author Contributions

Conceptualization: JDL DM SBZ.

Data curation: JM SBZ.

Formal analysis: JM SBZ.

Funding acquisition: JDL DM SBZ.

Investigation: JM SBZ US.

Methodology: JDL DM SBZ.

Project administration: TD JDL DM.

Resources: JDL DM.

Supervision: TD JDL AST DPW DM SBZ.

Visualization: JM SBZ.

Writing – original draft: JM.

Writing – review & editing: US TD JDL AST DPW DM SBZ.

References

- Rowell LB. Human cardiovascular adjustments to exercise and thermal stress. *Physiol Rev* 1974; 54:75–159. PMID: [4587247](#)
- Hooker SP, Sui X, Colabianchi N, Vena J, Laditka J, LaMonte MJ, et al. Cardiorespiratory fitness as a predictor of fatal and nonfatal stroke in asymptomatic women and men. *Stroke* 2008; 39:2950–2957. doi: [10.1161/STROKEAHA.107.495275](#) PMID: [18688008](#)
- LaMonte MJ, Barlow CE, Jurca R, Kampert JB, Church TS, Blair SN. Cardiorespiratory fitness is inversely associated with the incidence of metabolic syndrome: a prospective study of men and women. *Circulation* 2005; 112:505–512. doi: [10.1161/CIRCULATIONAHA.104.503805](#) PMID: [16009797](#)
- Katzmarzyk PT, Church TS, Craig CL, Bouchard C. Sitting time and mortality from all causes, cardiovascular disease, and cancer. *Med Sci Sports Exerc* 2009; 41:998–1005. doi: [10.1249/MSS.0b013e3181930355](#) PMID: [19346988](#)
- Zhang T, Zhang CF, Jin F, Wang L. Association between genetic factor and physical performance. *Yi Chuan* 2004; 26:219–226. PMID: [15662737](#)
- Jackson AS, Beard EF, Wier LT, Ross RM, Stuteville JE, Blair SN. Changes in aerobic power of men, ages 25–70 yr. *Med Sci Sports Exerc* 1995; 27:113–120. PMID: [7898326](#)
- Jackson AS, Wier LT, Ayers GW, Beard EF, Stuteville JE, Blair SN. Changes in aerobic power of women, ages 20–64 yr. *Med Sci Sports Exerc* 1996; 28:884–891. PMID: [8832543](#)
- Kenney WL, Wilmore JH, Costill DL. *Physiology of Sport and Exercise*. 5th ed. Champaign, IL: Human Kinetics; 2011. pp. 265–270.
- McArdle WD, Katch FI, Katch VL. *Exercise physiology: nutrition, energy, and human performance*. 8th ed. Baltimore: Wolters Kluwer Health; 2015. pp. 165–167.
- World Health Organisation. Global recommendations on physical activity for health. Genf. 2010. http://apps.who.int/iris/bitstream/10665/44399/1/9789241599979_eng.pdf. Accessed 7 August 2015.
- Ong TC, Sothy SP. A comparative study of the maximum oxygen uptake of regularly exercising and non-exercising health adult men in sedentary occupations. *Occup Med (Lond)* 1992; 42:120–124.
- Jang TW, Park SG, Kim HR, Kim JM, Hong YS, Kim BG. Estimation of maximal oxygen uptake without exercise testing in Korean healthy adult workers. *Tohoku J Exp Med* 2012; 227:313–319. PMID: [22850649](#)
- Perkio-Makela M. Influence of exercise-focused group activities on the physical activity, functional capacity, and work ability of female farmers—a three-year follow-up. *Int J Occup Saf Ergon* 1999; 5:381–394. doi: [10.1080/10803548.1999.11076427](#) PMID: [10602655](#)
- Castillo-Retamal M, Hinckson EA. Measuring physical activity and sedentary behaviour at work: a review. *Work* 2011; 40:345–357. doi: [10.3233/WOR-2011-1246](#) PMID: [22130052](#)
- Elliott SA, Baxter KA, Davies PS, Truby H. Accuracy of self-reported physical activity levels in obese adolescents. 2014. *J Nutr Metab*.
- Prince SA, Adamo KB, Hamel ME, Hardt J, Connor Gorber S, Tremblay M. A comparison of direct versus self-report measures for assessing physical activity in adults: a systematic review. *Int J Behav Nutr Phys Act* 2008; 5:56. doi: [10.1186/1479-5868-5-56](#) PMID: [18990237](#)
- Westerterp KR. Assessment of physical activity: a critical appraisal. *Eur J Appl Physiol* 2009; 105:823–828. doi: [10.1007/s00421-009-1000-2](#) PMID: [19205725](#)

18. Dishman RK. The measurement conundrum in exercise adherence research. *Med Sci Sports Exerc* 1994; 26:1382–1390. PMID: [7837959](#)
19. Freedson P, Bowles HR, Troiano R, Haskell W. Assessment of physical activity using wearable monitors: recommendations for monitor calibration and use in the field. *Med Sci Sports Exerc* 2012; 44:1–4.
20. Johannsen DL, Calabro MA, Stewart J, Franke W, Rood JC, Welk GJ. Accuracy of armband monitors for measuring daily energy expenditure in healthy adults. *Med Sci Sports Exerc* 2010; 42:2134–2140. doi: [10.1249/MSS.0b013e3181e0b3ff](#) PMID: [20386334](#)
21. World Health Organisation. Obesity and overweight. 2015. <http://www.who.int/mediacentre/factsheets/fs311/en/>. Accessed 7 August 2015.
22. Elias P. Occupational Classification (ISCO-88): Concepts, Methods, Reliability, Validity and Cross-National Comparability. 1997.
23. Steele R, Mummery K. Occupational physical activity across occupational categories. *J Sci Med Sport* 2003; 6:398–407. PMID: [14723390](#)
24. Leger L, Gadoury C. Validity of the 20 m shuttle run test with 1 min stages to predict VO₂max in adults. *Can J Sport Sci* 1989; 14:21–26. PMID: [2924218](#)
25. Cooper CB, Storer TW. Exercise Testing and Interpretation: A Practical Approach. 1st ed. Cambridge, United Kingdom: Cambridge University Press; 2001. pp. 57–59.
26. Uth N, Sorensen H, Overgaard K, Pedersen PK. Estimation of VO₂max from the ratio between HRmax and HRrest—the Heart Rate Ratio Method. *Eur J Appl Physiol* 2004; 91:111–115. doi: [10.1007/s00421-003-0988-y](#) PMID: [14624296](#)
27. Scheers T, Philippaerts R, Lefevre J. Variability in physical activity patterns as measured by the Sense-Wear Armband: how many days are needed? *Eur J Appl Physiol* 2012; 112:1653–1662. doi: [10.1007/s00421-011-2131-9](#) PMID: [21874552](#)
28. Pitta F, Takaki MY, Oliveira NH, Sant'anna TJ, Fontana AD, Kovelis D, et al. Relationship between pulmonary function and physical activity in daily life in patients with COPD. *Respir Med* 2008; 102:1203–1207. doi: [10.1016/j.rmed.2008.03.004](#) PMID: [18573647](#)
29. Garber CE, Blissmer B, Deschenes MR, Franklin BA, Lamonte MJ, Lee IM, et al. American College of Sports Medicine position stand. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal, and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011; 43:1334–1359. doi: [10.1249/MSS.0b013e318213febf](#) PMID: [21694556](#)
30. Committee AGA. Activity Guidelines Advisory Committee report, 2008. To the Secretary of Health and Human Services. Part A: executive summary. *Nutr Rev* 2008; 67:114–120.
31. Wultsch G, Rinnerhofer S, Tschakert G, Hofmann P. Governmental regulations for early retirement by means of energy expenditure cut offs. *Scand J Work Environ Health* 2012; 38:370–379. doi: [10.5271/sjweh.3195](#) PMID: [21918790](#)
32. Suminski RR, Wier LT, Poston W, Arenare B, Randles A, Jackson AS. The effect of habitual smoking on measured and predicted VO₂(max). *J Phys Act Health* 2009; 6:667–673. PMID: [19953845](#)
33. Swain DP, Franklin BA. VO₂(2) reserve and the minimal intensity for improving cardiorespiratory fitness. *Med Sci Sports Exerc* 2002; 34:152–157. PMID: [11782661](#)
34. Hammermeister J, Page RM, Dolny D, Burnham T. Occupational physical activity as an indicator of health and fitness. *Percept Mot Skills* 2001; 92:121–127. doi: [10.2466/pms.2001.92.1.121](#) PMID: [11322575](#)
35. Hirai T, Kusaka Y, Suganuma N, Seo A, Tobita Y. Physical work load affects the maximum oxygen uptake. *Ind Health* 2006; 44:250–257. PMID: [16715999](#)
36. Miller R, Brown W. Steps and sitting in a working population. *Int J Behav Med* 2004; 11:219–224. doi: [10.1207/s15327558ijbm1104_5](#) PMID: [15657022](#)
37. Swiss Federal Statistical Office. Swiss Labour Force Survey. <https://www.bfs.admin.ch/bfs/de/home/statistiken/arbeit-erwerb.assetdetail.255696.html>. Accessed 15 August 2015.
38. Bradshaw DI, George JD, Hyde A, LaMonte MJ, Vehrs PR, Hager RL, et al. An accurate VO₂max non-exercise regression model for 18–65-year-old adults. *Res Q Exerc Sport* 2005; 76:426–432. doi: [10.1080/02701367.2005.10599315](#) PMID: [16739680](#)
39. Mutikainen S, Helander E, Pietila J, Korhonen I, Kujala UM. Objectively measured physical activity in Finnish employees: a cross-sectional study. *BMJ Open* 2014; 4:e005927. doi: [10.1136/bmjopen-2014-005927](#) PMID: [25500160](#)
40. Swiss Federal Statistical Office. Physical Activity. <https://www.bfs.admin.ch/bfs/de/home/statistiken/gesundheit/determinanten/koerperliche-aktivitaet.html>. Accessed 17 August 2015.

41. Ainsworth BE, Haskell WL, Herrmann SD, Meckes N, Bassett DR Jr., Tudor-Locke C, et al. 2011 Compendium of Physical Activities: a second update of codes and MET values. *Med Sci Sports Exerc* 2011; 43:1575–1581. doi: [10.1249/MSS.0b013e31821ece12](https://doi.org/10.1249/MSS.0b013e31821ece12) PMID: [21681120](https://pubmed.ncbi.nlm.nih.gov/21681120/)
42. Drenowatz C, Eisenmann JC. Validation of the SenseWear Armband at high intensity exercise. *Eur J Appl Physiol* 2011; 111:883–887. doi: [10.1007/s00421-010-1695-0](https://doi.org/10.1007/s00421-010-1695-0) PMID: [20972880](https://pubmed.ncbi.nlm.nih.gov/20972880/)
43. Wetten AA, Batterham M, Tan SY, Tapsell L. Relative validity of 3 accelerometer models for estimating energy expenditure during light activity. *J Phys Act Health* 2014; 11:638–647. doi: [10.1123/jpah.2011-0167](https://doi.org/10.1123/jpah.2011-0167) PMID: [23417054](https://pubmed.ncbi.nlm.nih.gov/23417054/)
44. Wasserman K. Diagnosing cardiovascular and lung pathophysiology from exercise gas exchange. *Chest* 1997; 112:1091–1101. PMID: [9377922](https://pubmed.ncbi.nlm.nih.gov/9377922/)
45. Ramsbottom R, Brewer J, Williams C. A progressive shuttle run test to estimate maximal oxygen uptake. *Br J Sports Med* 1988; 22:141–144. PMID: [3228681](https://pubmed.ncbi.nlm.nih.gov/3228681/)
46. Paradisis GP, Zacharogiannis E, Mandila D, Smirtiotou A, Argeitaki P, Cooke CB. Multi-Stage 20-m Shuttle Run Fitness Test, Maximal Oxygen Uptake and Velocity at Maximal Oxygen Uptake. *J Hum Kinet* 2014; 41:81–87. doi: [10.2478/hukin-2014-0035](https://doi.org/10.2478/hukin-2014-0035) PMID: [25114734](https://pubmed.ncbi.nlm.nih.gov/25114734/)